



Solve each problem.

**Answers**

1) Which table of values can be defined by the function:  $y = x + 4$

A.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-1</td><td>-11</td></tr><tr><td>0</td><td>-7</td></tr><tr><td>1</td><td>-3</td></tr><tr><td>2</td><td>1</td></tr></tbody></table>	x	y	-1	-11	0	-7	1	-3	2	1	B.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>-3</td></tr><tr><td>-2</td><td>-2</td></tr><tr><td>-1</td><td>-1</td></tr><tr><td>0</td><td>0</td></tr></tbody></table>	x	y	-3	-3	-2	-2	-1	-1	0	0	C.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-2</td><td>-1</td></tr><tr><td>-1</td><td>3</td></tr><tr><td>0</td><td>7</td></tr><tr><td>1</td><td>11</td></tr></tbody></table>	x	y	-2	-1	-1	3	0	7	1	11	D.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>1</td></tr><tr><td>-2</td><td>2</td></tr><tr><td>-1</td><td>3</td></tr><tr><td>2</td><td>6</td></tr></tbody></table>	x	y	-3	1	-2	2	-1	3	2	6
x	y																																														
-1	-11																																														
0	-7																																														
1	-3																																														
2	1																																														
x	y																																														
-3	-3																																														
-2	-2																																														
-1	-1																																														
0	0																																														
x	y																																														
-2	-1																																														
-1	3																																														
0	7																																														
1	11																																														
x	y																																														
-3	1																																														
-2	2																																														
-1	3																																														
2	6																																														

1. \_\_\_\_\_

2. \_\_\_\_\_

3. \_\_\_\_\_

4. \_\_\_\_\_

5. \_\_\_\_\_

2) Which table of values can be defined by the function:  $y = x \times (-8)$

A.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-2</td><td>-2</td></tr><tr><td>-1</td><td>-1</td></tr><tr><td>1</td><td>1</td></tr><tr><td>2</td><td>2</td></tr></tbody></table>	x	y	-2	-2	-1	-1	1	1	2	2	B.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>24</td></tr><tr><td>-2</td><td>16</td></tr><tr><td>1</td><td>-8</td></tr><tr><td>2</td><td>-16</td></tr></tbody></table>	x	y	-3	24	-2	16	1	-8	2	-16	C.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>-216</td></tr><tr><td>-1</td><td>-72</td></tr><tr><td>0</td><td>0</td></tr><tr><td>1</td><td>72</td></tr></tbody></table>	x	y	-3	-216	-1	-72	0	0	1	72	D.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-2</td><td>6</td></tr><tr><td>0</td><td>8</td></tr><tr><td>3</td><td>11</td></tr><tr><td>4</td><td>12</td></tr></tbody></table>	x	y	-2	6	0	8	3	11	4	12
x	y																																														
-2	-2																																														
-1	-1																																														
1	1																																														
2	2																																														
x	y																																														
-3	24																																														
-2	16																																														
1	-8																																														
2	-16																																														
x	y																																														
-3	-216																																														
-1	-72																																														
0	0																																														
1	72																																														
x	y																																														
-2	6																																														
0	8																																														
3	11																																														
4	12																																														

3) Which table of values can be defined by the function:  $y = 9x \times 7$

A.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>-13</td></tr><tr><td>-3</td><td>-12</td></tr><tr><td>-2</td><td>-11</td></tr><tr><td>1</td><td>-8</td></tr></tbody></table>	x	y	-4	-13	-3	-12	-2	-11	1	-8	B.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>-189</td></tr><tr><td>-2</td><td>-126</td></tr><tr><td>-1</td><td>-63</td></tr><tr><td>1</td><td>63</td></tr></tbody></table>	x	y	-3	-189	-2	-126	-1	-63	1	63	C.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>-3</td></tr><tr><td>-1</td><td>-1</td></tr><tr><td>1</td><td>1</td></tr><tr><td>3</td><td>3</td></tr></tbody></table>	x	y	-3	-3	-1	-1	1	1	3	3	D.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>27</td></tr><tr><td>-2</td><td>18</td></tr><tr><td>1</td><td>-9</td></tr><tr><td>3</td><td>-27</td></tr></tbody></table>	x	y	-3	27	-2	18	1	-9	3	-27
x	y																																														
-4	-13																																														
-3	-12																																														
-2	-11																																														
1	-8																																														
x	y																																														
-3	-189																																														
-2	-126																																														
-1	-63																																														
1	63																																														
x	y																																														
-3	-3																																														
-1	-1																																														
1	1																																														
3	3																																														
x	y																																														
-3	27																																														
-2	18																																														
1	-9																																														
3	-27																																														

4) Which table of values can be defined by the function:  $y = 3x + 6$

A.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-1</td><td>3</td></tr><tr><td>0</td><td>6</td></tr><tr><td>1</td><td>9</td></tr><tr><td>2</td><td>12</td></tr></tbody></table>	x	y	-1	3	0	6	1	9	2	12	B.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-1</td><td>2</td></tr><tr><td>0</td><td>3</td></tr><tr><td>1</td><td>4</td></tr><tr><td>3</td><td>6</td></tr></tbody></table>	x	y	-1	2	0	3	1	4	3	6	C.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>12</td></tr><tr><td>-1</td><td>3</td></tr><tr><td>2</td><td>-6</td></tr><tr><td>3</td><td>-9</td></tr></tbody></table>	x	y	-4	12	-1	3	2	-6	3	-9	D.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>-12</td></tr><tr><td>0</td><td>0</td></tr><tr><td>1</td><td>3</td></tr><tr><td>3</td><td>9</td></tr></tbody></table>	x	y	-4	-12	0	0	1	3	3	9
x	y																																														
-1	3																																														
0	6																																														
1	9																																														
2	12																																														
x	y																																														
-1	2																																														
0	3																																														
1	4																																														
3	6																																														
x	y																																														
-4	12																																														
-1	3																																														
2	-6																																														
3	-9																																														
x	y																																														
-4	-12																																														
0	0																																														
1	3																																														
3	9																																														

5) Which table of values can be defined by the function:  $y = 8x \div 8$

A.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>-288</td></tr><tr><td>-2</td><td>-144</td></tr><tr><td>-1</td><td>-72</td></tr><tr><td>0</td><td>0</td></tr></tbody></table>	x	y	-4	-288	-2	-144	-1	-72	0	0	B.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-3</td><td>-19</td></tr><tr><td>-1</td><td>-1</td></tr><tr><td>2</td><td>26</td></tr><tr><td>3</td><td>35</td></tr></tbody></table>	x	y	-3	-19	-1	-1	2	26	3	35	C.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>-13</td></tr><tr><td>-3</td><td>-12</td></tr><tr><td>0</td><td>-9</td></tr><tr><td>1</td><td>-8</td></tr></tbody></table>	x	y	-4	-13	-3	-12	0	-9	1	-8	D.	<table border="1"><thead><tr><th>x</th><th>y</th></tr></thead><tbody><tr><td>-4</td><td>-4</td></tr><tr><td>-2</td><td>-2</td></tr><tr><td>-1</td><td>-1</td></tr><tr><td>0</td><td>0</td></tr></tbody></table>	x	y	-4	-4	-2	-2	-1	-1	0	0
x	y																																														
-4	-288																																														
-2	-144																																														
-1	-72																																														
0	0																																														
x	y																																														
-3	-19																																														
-1	-1																																														
2	26																																														
3	35																																														
x	y																																														
-4	-13																																														
-3	-12																																														
0	-9																																														
1	-8																																														
x	y																																														
-4	-4																																														
-2	-2																																														
-1	-1																																														
0	0																																														



Solve each problem.

1) Which table of values can be defined by the function:  $y = x + 4$

A. 

x	y
-1	-11
0	-7
1	-3
2	1

B. 

x	y
-3	-3
-2	-2
-1	-1
0	0

C. 

x	y
-2	-1
-1	3
0	7
1	11

D. 

x	y
-3	1
-2	2
-1	3
2	6

2) Which table of values can be defined by the function:  $y = x \times (-8)$

A. 

x	y
-2	-2
-1	-1
1	1
2	2

B. 

x	y
-3	24
-2	16
1	-8
2	-16

C. 

x	y
-3	-216
-1	-72
0	0
1	72

D. 

x	y
-2	6
0	8
3	11
4	12

3) Which table of values can be defined by the function:  $y = 9x \times 7$

A. 

x	y
-4	-13
-3	-12
-2	-11
1	-8

B. 

x	y
-3	-189
-2	-126
-1	-63
1	63

C. 

x	y
-3	-3
-1	-1
1	1
3	3

D. 

x	y
-3	27
-2	18
1	-9
3	-27

4) Which table of values can be defined by the function:  $y = 3x + 6$

A. 

x	y
-1	3
0	6
1	9
2	12

B. 

x	y
-1	2
0	3
1	4
3	6

C. 

x	y
-4	12
-1	3
2	-6
3	-9

D. 

x	y
-4	-12
0	0
1	3
3	9

5) Which table of values can be defined by the function:  $y = 8x \div 8$

A. 

x	y
-4	-288
-2	-144
-1	-72
0	0

B. 

x	y
-3	-19
-1	-1
2	26
3	35

C. 

x	y
-4	-13
-3	-12
0	-9
1	-8

D. 

x	y
-4	-4
-2	-2
-1	-1
0	0

Answers

1. **D**

2. **B**

3. **B**

4. **A**

5. **D**